



The CRASH Code

**NIF User Group Meeting
February, 2012**

Ken Powell, R. Paul Drake, James Holloway, Quentin Stout, Marv Adams,
Gabor Toth, Bart van der Holst, Igor Sokolov, Ben Torralva, Eric Myra,
Bruce Fryxell, Mike Grosskopf and Erica Rutter and the rest of the CRASH
Team




CRASH is a radiation-hydrodynamic code with a laser package

- 1D, 2D or 3D
- Dynamic adaptive AMR
- Level set interfaces
- Self-consistent EOS and opacities
- Multigroup-diffusion radiation transport
- Electron physics and flux-limited electron heat conduction
- Laser package
 - 3D ray tracing for 2D or 3D runs


3D Nozzle to Ellipse @ 13 ns




Material & AMR



Log Density



Log Electron Temperature




Log Ion Temperature

CRASH code: Van der Holst et al, Ap.J.S. 2011




The genesis of CRASH was space-weather modeling

- We developed a practical high-resolution Godunov scheme for multi-dimensional MHD
- We built an efficient solution-adaptive parallel MHD solver (BATS-R-US)
- We tied a number of physics modules (some built on BATS-R-US) together to form the Space Weather Modeling Framework (SWMF)






The disparate scales of space-weather drove our need for solution-adaptive parallel code

- Temporal scale range:
 - $\sim 2^{28} \approx 2.5 \times 10^8$
- Linear spatial scale range:
 - $\sim 2^{28} \approx 2.5 \times 10^8$




BATSRUS can model many plasma processes

- Compressible fluid dynamics
- Ideal MHD
- Resistive MHD
- Hall MHD
- Semi-relativistic MHD
- Physics-based energy transport
 - Heat conduction
 - Wave energy transport
- Multi-fluid MHD
 - Each ionic species has its own continuity, momentum and energy equation
 - Electron momentum equation is replaced by Ohm's law.





The Space-Weather Modeling Framework comprises many modules



SWMF has been extensively validated

- In late October and early November 2003 a series of some of the most powerful solar eruptions ever registered occurred.
- The “Halloween storm” simulation provided a unique opportunity for code/observation comparison



CRASH is an extension of BATS-R-US to the high-energy-density domain

- Builds on the solution-adaptive, parallel framework of BATS-R-US
- Adds new physics
 - Radiation transport
 - Electron physics and flux-limited electron heat conduction
 - Laser package
 - Tabular and self-consistent EOS and opacities
- Can tap into physics in BATS-R-US (e.g. MHD)



We model laser energy transport using a parallel ray-tracing algorithm for AMR grids

Rays are traced by solving



$$\frac{d\mathbf{r}^2}{ds^2} = \frac{d\mathbf{r}}{ds} \times \left(\frac{\nabla n}{n} \times \frac{d\mathbf{r}}{ds} \right)$$

where \mathbf{r} is the ray direction, s is the distance along the ray, and n is the index of refraction

∇n is determined from the plasma density distribution

Laser energy is absorbed by electron-ion collisions

The laser energy is smoothly deposited in the plasma by distributing it among the nearest cells



CRASH Radhydro Code: Hydro and Electron Physics

$$\frac{\partial}{\partial t} \left\{ \begin{array}{c} \rho \\ \rho \mathbf{u} \\ \mathcal{E} + \frac{1}{2} \rho \mathbf{u} \cdot \mathbf{u} \\ \mathcal{E}_e \end{array} \right\} + \nabla \cdot \left\{ \begin{array}{c} \rho \mathbf{u} \\ \rho \mathbf{u} \mathbf{u} + p \mathbf{I} \\ \mathbf{u} \left(\frac{1}{2} \rho \mathbf{u} \cdot \mathbf{u} + \mathcal{E} + p \right) \\ \mathbf{u} \mathcal{E}_e \end{array} \right\} = \mathbf{S}$$

$$\mathbf{S} = \left\{ \begin{array}{l} \text{electron heat conduction} \\ -p_e \nabla \cdot \mathbf{u} + \nabla \cdot C_e \nabla T_e + \frac{\rho k_B (T_i - T_e)}{M_p A \tau_{ei}} - (S_{re} - \mathbf{S}_{rm} \cdot \mathbf{u}) + S_L \end{array} \right\}$$

Diagram illustrating the components of the source term \mathbf{S} in the CRASH Radhydro Code:

- laser energy deposition**: Points to S_L .
- radiation/electron momentum exchange**: Points to $-\mathbf{S}_{rm}$.
- electron heat conduction**: Points to $\nabla \cdot C_e \nabla T_e$.
- Collisional exchange**: Points to $\frac{\rho k_B (T_i - T_e)}{M_p A \tau_{ei}}$.
- radiation/electron energy exchange**: Points to S_{re} .
- Compression work**: Points to $-p_e \nabla \cdot \mathbf{u}$.

CRASH Radhydro Code: Multigroup diffusion


- Radiation transport equation reduces to a system of equations for spectral energy density of groups.
- Diffusion is flux-limited
- For the g^{th} group:

$$\frac{\partial E_g}{\partial t} + \underbrace{\nabla \cdot (E_g \mathbf{u})}_{\text{advection}} - \underbrace{p_g \nabla \cdot \mathbf{u}}_{\text{compression work}} - \underbrace{\frac{\nabla \cdot \mathbf{u}}{\Delta(\log \varepsilon)} \Delta(p_g)}_{\text{photon energy shift}} = \text{diffusion} + \text{emission} - \text{absorption}$$

$$\text{diffusion} = \nabla \cdot (D_g \nabla E_g) \quad \text{emission-absorption} = c \chi_{abs_g} (B_g - E_g)$$

$$\Delta(\cdot) = (\cdot)_{g+\frac{1}{2}} - (\cdot)_{g-\frac{1}{2}}$$

Overview of Solver Approach




- Self-similar block-based adaptive grid
- Finite-volume scheme, approximate Riemann solver for flux function, limited linear interpolation
- Level-set equations used to evolve material interfaces; each cell treated as single-material cell
- Mixed Implicit/Explicit update
 - Hydro and electron equations
 - Advection, compression and pressure force updated explicitly
 - Exchange terms and electron heat conduction treated implicitly
 - Radtran
 - Advection of radiation energy, compression work and photon shift are evaluated explicitly
 - Diffusion and emission-absorption are evaluated implicitly
 - Implicit scheme is a preconditioned Newton-Krylov-Schwarz scheme

We extensively test our code


- New program units implemented with **unit tests**
 - Nightly execution of many unit tests for CRASH and its parent code
- New features implemented with **verification tests**
 - Daily verification & **full system tests** are run on a 16-core Mac.
 - Tests cover all aspects of the new feature, including restart, using grid convergence studies and model-model comparison.
- Compatibility & reproducibility checked with **functionality test suite**
 - Nightly runs. 9 different platforms/compilers on 1 to 4 cores: tests portability
- **Parallel Scaling Tests**
 - Weekly scaling test on 128 and 256 cores of hera.
 - Reveals software and hardware issues, and confirms that results are independent of the number of cores.

Multiple classes of tests are in our suite


HEAT CONDUCTION



RADIATION TRANSPORT




FULL SYSTEM




- Hydrodynamics
- Radiation transport
- Radiation hydrodynamics
- Heat conduction
- Simulated radiography
- Material properties (EOS and opacities)
- Unit tests
- Full-system tests


HYDRODYNAMICS



RADIATION HYDRODYNAMICS




SIMULATED RADIOGRAPHY




The hydro portion of the code scales well to tens of thousands of cores

(CRASH hydro Weak Scaling on BG/L)






Radiation-hydrodynamics scales well to ~ 1000 cores

CRASH rad-hydro strong scaling on Hera and Pleiades




The CRASH project itself is focused on radiative shocks


- 1 ns, 4 kJ laser irradiates 20 μm Be disk
- Drives shock into Xe-filled tube at 1.1 atm.
- Radiative precursor heats wall of tube, leading to ablation
- Complex interaction among laser-driven shock, ablation-driven shock, and Xe-Be interface




Full system with CRASH laser package




We can now simulate the Year-5 Experiment




CRASH simulation of Y5 experiment



CRASH simulation of Y5 experiment




radiograph




time = 1.8 ns


radiograph



time = 1.8 ns




CRASH has been used to model several HED experiments




Rayleigh-Taylor growth in a diverging system

High Drive:
310 eV Tr source


Low Drive
207 eV Tr source



Rayleigh-Taylor growth in the presence of a radiative shock



Ablative flow of laser driven foil for collisionless shock experiments



Creation of plasma jets using laser irradiation of conical foils

See Poster by M J Grosskopf for more details as well as information on other applications being modeled with the CRASH code



Concluding Remarks

- The CRASH code is now useful for applications
- We follow good practices on code development and verification
- We have simulated the experiments for the CRASH project
- We have simulated several other HED experiments
- The code is publicly available
 - But realistically requires knowledge and experience to run
 - We welcome visitors seeking these



Acknowledgments

The Center for Radiative Shock Hydrodynamics, which developed the CRASH code, is funded by the Predictive Sciences Academic Alliances Program in NNSA-ASC via grant DEFC52- 08NA28616.

Related experimental work, often now using CRASH, is in the Center for Laser Experimental Astrophysics Research, funded by the NNSA-DS and SC-OFES Joint Program in High-Energy-Density Laboratory Plasmas, grant number DE-FG52-09NA29548, by the National Laser User Facility Program, grant number DE-FG52-09NA29034, and by other sponsors.

